

## POSTOPERATIVE DISCHARGE INSTRUCTIONS MENISCAL REPAIR

Benjamin D. Rubin, M.D.  
714 937-2120

### DAY OF SURGERY

- **Diet** - start with clear liquids and gradually increase to a regular diet as tolerated. It is not uncommon to have some nausea or even vomiting after general anesthesia.

- **Ice** – If using the gel packs provided by the surgery center, keep them on until they thaw and then change to the alternate packs that are in the freezer for the first 2 days. After the dressing is changed, ice 20 minutes on/20 minutes off.

***Do not apply the ice directly to your skin as this can cause a severe burn and wound complications.***

Always keep a pad between the ice pack and your skin. If you have been provided with a cold unit, keep this in place. The cold unit can be used continuously since it is temperature controlled at a temperature setting between 1 & 2 on the unit.

- **Pain Medication** – On the day of the surgery if the local anesthetic is working and there is no significant pain, then before going to bed take a pain pill and place another on the nightstand with some water to take during the night if you are awakened with pain.

- **Clot (DVT) Prevention**—Take aspirin 325 mg daily for 2 weeks post-op and perform ankle pumps (as instructed after surgery) to decrease the incidence of developing a blood clot.

Take the prescribed pain medication at the onset of pain as needed. Do not try to tough it out but do not take it unnecessarily. If you are not having much in the way of pain you can take Ibuprofen, Advil or Aleve instead. Please take it with food to avoid stomach upset. If the medication gives inadequate relief or if you develop a reaction to the pain medication, stop taking it and call my office preferably during office hours.

- **Bandages** – The bandage applied to your knee may become moist or bloodstained, however this is normal and results from fluid placed in the joint during surgery. If your bandage becomes saturated with bright red blood, please call me. Keep the bandages dry.

- **Activity** - Ice and elevate your leg preferably above the level of your heart (on 2 pillows). Do not overdo your activity as your knee will probably be numb from the local anesthetic that was injected at the end of the procedure. This may last from 6 to 12 hours.

### 2. DAY AFTER SURGERY UNTIL FIRST POSTOPERATIVE OFFICE VISIT

- **Ice** – Continue to ice as much as possible to control the pain and swelling. If using the gel packs provided by the surgery center, keep them on until they thaw and then change to the alternate packs that are in the freezer for the first 2 days. After the dressing is changed, ice 20 minutes on/20 minutes off.

- **Activity** – Partial weight bear with the crutches in TROM brace locked out straight for 2 weeks. When seated you can remove the brace or use the drop lock hinges on the side of the brace to flex to 90° but not beyond.

- **Bandages** – You can remove the outer bandages and shower on the second day after surgery (48 hours). **Do not remove the tapes over your incisions.** Do not submerge your knee in a bath or spa. It is normal to have some swelling, drainage and minor redness at the incisions. It is common to have swelling of the foot and ankle.

- **Pain medication** – As needed. Please anticipate your needs for medication. Prescriptions are not renewed on weekends.

**PRECAUTION:** If you develop a fever of 101 degrees or greater, if the pain you feel is much greater than that expected, or if you develop severe redness or swelling, please call my office.